arXiv:1108.1171v2 [math.NT] 25 Mar 2012

PROOF OF A CONGRUENCE FOR HARMONIC NUMBERS CONJECTURED BY Z.-W. SUN

ROMEO MEŠTROVIĆ

Abstract. For a positive integer n let $H_n=\sum_{k=1}^{n}1/k$ be the nth harmonic number. In this note we prove that for any prime $p\ge 7$,

$$\sum_{k=1}^{p-1}\frac{H_k^2}{k^2}\equiv\frac{4}{5}pB_{p-5}\pmod{p^2},$$

which confirms the conjecture recently proposed by Z.-W. Sun. Furthermore, we also prove two similar congruences modulo p^2.

1. **The Main Result and Its Proof**

For a positive integer n, the nth *harmonic number* H_n is defined as

$$H_n=\sum_{k=1}^{n}\frac{1}{k}$$

(in addition, we define $H_0=0$).

Recently, Zhi-Wei Sun [5] obtained basic congruences modulo a prime $p\ge 5$ for several sums of terms involving harmonic numbers. In particular, Sun established $\sum_{k=1}^{p-1}H_k^r\,(\mathrm{mod}\,p^{4-r})$ for $r=1,2,3$. Further generalizations of these congruences have been recently obtained by Tauraso in [8]. Notice also that Sun [5, Theorem 1.1] proved that for any prime $p>5$

$$\sum_{k=1}^{p-1}\frac{H_k^2}{k^2}\equiv 0\pmod{p},\tag{1.1}$$

and conjectured [5, the second part of Conjecture 1.1] (also see [6, Conjecture A37 (v), p. 31]) that for such a prime p,

$$\sum_{k=1}^{p-1}\frac{H_k^2}{k^2}\equiv\frac{4}{5}pB_{p-5}\pmod{p^2},\tag{1.2}$$

where $B_0,B_1,B_2,\dots,$ are Bernoulli numbers given by

$$B_0=1\quad\text{and}\quad\sum_{k=0}^{n}\binom{n+1}{k}B_k=0\ (n=1,2,3,\dots).$$

Recall also that the first part of Conjecture 1.1 in [5] was recently confirmed by Z.-W. Sun and L. L. Zhao in [7, the congruence (1.1) of Theorem 1.1]; for another proof see [4, the congruence (7) of Corollary 1.2].

2010 *Mathematics Subject Classification.* Primary 11B75; Secondary 11A07, 11B68, 05A10.

Key words and phrases. Harmonic numbers, congruences, Bernoulli numbers.

In this note we prove the congruence (1.2) and two similar congruences contained in the following result.

Theorem 1.1. *Let $p>5$ be a prime. Then*

$$\sum_{k=1}^{p-1}\frac{H_k^2}{k^2}\equiv\frac{4}{5}pB_{p-5}\pmod{p^2},\tag{1.3}$$

$$\sum_{k=1}^{p-1}\frac{H_k^3}{k}\equiv\frac{3}{2}pB_{p-5}\pmod{p^2},\tag{1.4}$$

$$\sum_{k=1}^{p-1}\frac{H_k}{k^3}\equiv-\frac{1}{10}pB_{p-5}\pmod{p^2}.\tag{1.5}$$

The proof of Theorem 1.1 presented below is based on some congruences for multiple harmonic sums obtained by J. Zhao [9].

Proof of Theorem 1.1. For simplicity, here we denote

$$H(2,1,1)=\sum_{1\le i<j<k\le p-1}\frac{1}{i^2jk},\quad H(1,2,1)=\sum_{1\le i<j<k\le p-1}\frac{1}{ij^2k},$$

$$H(m,n)=\sum_{1\le i<j\le p-1}\frac{1}{i^mj^n},\ \text{for } m,n=1,2,\dots,\ H(n)=\sum_{k=1}^{p-1}\frac{1}{k^n}\ \text{for } n=1,2,\dots.$$

Proof of (1.3). By *Wolstenholme theorem* (see e.g., [1, Theorem 1] or [3, p. 255]), for any prime $p>3$, $H(1):=\sum_{k=1}^{p-1}1/k\equiv 0\,(\mathrm{mod}\,p^2)$, or equivalently, for each $j=1,2,\dots,p-2$ the following congruence holds

$$\frac{1}{j+1}+\frac{1}{j+2}+\cdots+\frac{1}{p-1}\equiv-\left(1+\frac{1}{2}+\cdots+\frac{1}{j-1}+\frac{1}{j}\right)\pmod{p^2}.\tag{1.6}$$

Applying the congruence (1.6), we find that

$$\begin{aligned}
H(1,2,1)&=\sum_{1\le i<j<k\le p-1}\frac{1}{ij^2k}=\sum_{j=2}^{p-1}\frac{1}{j^2}\sum_{i=1}^{j-1}\frac{1}{i}\sum_{k=j+1}^{p-1}\frac{1}{k}\\
&=\sum_{j=2}^{p-1}\frac{1}{j^2}\left(1+\frac{1}{2}+\cdots+\frac{1}{j-1}\right)\left(\frac{1}{j+1}+\frac{1}{j+2}+\cdots+\frac{1}{p-1}\right)\\
&\equiv\sum_{j=2}^{p-1}\frac{1}{j^2}\left(1+\frac{1}{2}+\cdots+\frac{1}{j-1}\right)\left(-\left(1+\frac{1}{2}+\cdots+\frac{1}{j}\right)\right)\pmod{p^2}\\
&=\sum_{j=1}^{p-1}\frac{1}{j^2}\left(H_j-\frac{1}{j}\right)(-H_j)=-\sum_{j=1}^{p-1}\frac{H_j^2}{j^2}+\sum_{j=1}^{p-1}\frac{H_j}{j^3}\\
&=-\sum_{j=1}^{p-1}\frac{H_j^2}{j^2}+\sum_{j=1}^{p-1}\frac{H_{j-1}+\frac{1}{j}}{j^3}=-\sum_{j=1}^{p-1}\frac{H_j^2}{j^2}+\sum_{1\le i<j\le p-1}\frac{1}{ij^3}+\sum_{j=1}^{p-1}\frac{1}{j^4}\\
&=-\sum_{j=1}^{p-1}\frac{H_j^2}{j^2}+H(1,3)+H(4)\qquad\pmod{p^2}.
\end{aligned}\tag{1.7}$$

From the shuffle relation (see e.g., [9, (3.6) on page 89])

$$H(1)H(3) = H(1,3) + H(3,1) + H(4)$$

and the fact that $p^2 \mid H(1)$ it follows that

$$H(1,3) + H(4) \equiv -H(3,1) \pmod{p^2}.$$

Substituting this into the right hand side of (1.7), we immediately obtain

$$H(1,2,1) + H(3,1) \equiv -\sum_{j=1}^{n} \frac{H_j^2}{j^2} \pmod{p^2}. \tag{1.8}$$

Further, by the shuffle relation

$$H(1)H(2,1) = 2H(2,1,1) + H(1,2,1) + H(3,1) + H(2,2)$$

again using the fact that $p^2 \mid H(1)$, we get

$$H(1,2,1) + H(3,1) \equiv -2H(2,1,1) - H(2,2) \pmod{p^2} \tag{1.9}$$

By the congruence (3.22) in [9, Proposition 2.7, p. 93], we have

$$H(2,1,1) \equiv \frac{3}{5} pB_{p-5} \pmod{p^2}, \tag{1.10}$$

and by [9, Theorem 3.2, p. 88] or [10],

$$H(2,2) \equiv -\frac{2}{5} pB_{p-5} \pmod{p^2}. \tag{1.11}$$

From (1.9), (1.10) and (1.11) we find that

$$H(1,2,1) + H(3,1) \equiv -\frac{4}{5} pB_{p-5} \pmod{p^2}, \tag{1.12}$$

which substituting into (1.8) implies the congruence (1.3).

Proof of (1.4). Since $H_k = H_{k-1} + 1/k$, for each $k = 1, 2, \ldots, p-1$ the binomial formula implies

$$\begin{aligned} H_k^4 - H_{k-1}^4 &= 4\frac{H_{k-1}^3}{k} + 6\frac{\left(H_k - \frac{1}{k}\right)^2}{k^2} + 4\frac{H_{k-1}}{k^3} + \frac{1}{k^4} \\ &= 4\frac{H_{k-1}^3}{k} + 6\left(\frac{H_k^2}{k^2} - 2\frac{H_{k-1} + \frac{1}{k}}{k^3} + \frac{1}{k^4}\right) + 4\frac{H_{k-1}}{k^3} + \frac{1}{k^4} \\ &= 4\frac{H_{k-1}^3}{k} + 6\frac{H_k^2}{k^2} - 8\frac{H_{k-1}}{k^3} - \frac{5}{k^4}. \end{aligned}$$

After summation of the above identity over k, and using the fact that by Wolstenholme theorem, $p^2 \mid H(1)$, we get

$$4\sum_{k=1}^{p-1} \frac{H_{k-1}^3}{k} + 6\sum_{k=1}^{p-1} \frac{H_k^2}{k^2} - 8\sum_{k=1}^{p-1} \frac{H_{k-1}}{k^3} - 5\sum_{k=1}^{p-1} \frac{1}{k^4} = H_{p-1}^4 \equiv 0 \pmod{p^2}. \tag{1.13}$$

Similarly, we have

$$\begin{aligned} \frac{H_{k-1}^3}{k} &= \frac{1}{k}\left(H_k - \frac{1}{k}\right)^3 = \frac{H_k^3}{k} - 3\frac{H_k^2}{k^2} + 3\frac{H_{k-1} + \frac{1}{k}}{k^3} - \frac{1}{k^4} \\ &= \frac{H_k^3}{k} - 3\frac{H_k^2}{k^2} + 3\frac{H_{k-1}}{k^3} + \frac{2}{k^4}, \end{aligned}$$

which gives

$$\sum_{k=1}^{p-1}\frac{H_{k-1}^3}{k}=\sum_{k=1}^{p-1}\frac{H_k^3}{k}-3\sum_{k=1}^{p-1}\frac{H_k^2}{k^2}+3\sum_{k=1}^{p-1}\frac{H_{k-1}}{k^3}+2\sum_{k=1}^{p-1}\frac{1}{k^4}. \tag{1.14}$$

Substituting the expression (1.14) into the first term of (1.13), we immediately obtain

$$4\sum_{k=1}^{p-1}\frac{H_k^3}{k}-6\sum_{k=1}^{p-1}\frac{H_k^2}{k^2}+4\sum_{k=1}^{p-1}\frac{H_{k-1}}{k^3}+3\sum_{k=1}^{p-1}\frac{1}{k^4}\equiv 0 \pmod{p^2}. \tag{1.15}$$

Further, since by [9, Theorem 3.2, p. 88] $H(1,3)\equiv -\frac{9}{10}pB_{p-5} \,(\mathrm{mod}\, p^2)$, we have

$$\sum_{k=1}^{p-1}\frac{H_{k-1}}{k^3}=\sum_{1\le i<k\le p-1}\frac{1}{ik^3}=H(1,3)\equiv -\frac{9}{10}pB_{p-5} \pmod{p^2}. \tag{1.16}$$

Inserting the congruence (1.11) and the well known congruence $H(2)\equiv 0 \,(\mathrm{mod}\, p)$ (see e.g., [2, Theorem 3]) into the shuffle relation $H(2)H(2)=2H(2,2)+H(4)$, we find that

$$\sum_{k=1}^{p-1}\frac{1}{k^4}=H(4)\equiv -2H(2,2)\equiv \frac{4}{5}pB_{p-5} \pmod{p^2}. \tag{1.17}$$

Substituting (1.3), (1.16) and (1.17) into (1.15), we obtain the congruence (1.4).

Proof of (1.5). The congruences (1.16) and (1.17) yield

$$\sum_{k=1}^{p-1}\frac{H_k}{k^3}=\sum_{k=1}^{p-1}\frac{H_{k-1}}{k^3}+\sum_{k=1}^{p-1}\frac{1}{k^4}=H(1,3)+H(4)\equiv -\frac{1}{10}pB_{p-5} \pmod{p^2},$$

as desired. This completes the proof of Theorem 1.1. □

References

[1] E. Alkan, Variations on Wolstenholme's theorem, *Amer. Math. Monthly* **101** (1994) 1001–1004.

[2] M. Bayat, A generalization of Wolstenholme's Theorem, *Amer. Math. Monthly* **104** (1997) 557–560.

[3] A. Granville, Arithmetic properties of binomial coefficients. I. Binomial coefficients modulo prime powers, in *Organic Mathematics* (*Burnaby, BC*, 1995), *CMS Conf. Proc.*, Vol. 20 (American Mathematical Society, Providence, RI, 1997), pp. 253–276,

[4] R. Meštrović, On the mod p^2 determination of $\sum_{k=1}^{p-1}H_k/(k\cdot 2^k)$: another proof of a conjecture by Sun, accepted for publication in *Publ. Math. Debrecen*, preprint `arXiv:1108.3197 [math.NT]` (2011).

[5] Z.-W. Sun, Arithmetic theory of harmonic numbers, *Proc. Amer. Math. Soc.* **140** (2012) 415–428.

[6] Z.-W. Sun, Open conjectures on congruences, preprint `arXiv:0911.5665v54 [math.NT]` (2011).

[7] Z.-W. Sun and L. L. Zhao, Arithmetic theory of harmonic numbers (II), preprint `arXiv:0911.4433v7 [math.NT]` (2011).

[8] R. Tauraso, New harmonic number identities with applications, *Séminaire Lotharingien de Combinatoire* **63** (2010) Article B63g.

[9] J. Zhao, Wolstenholme type theorem for multiple harmonic sums, *Int. J. Number Theory* **4** (2008) 73–106.

[10] X. Zhou and T. Cai, A generalization of a curious congruence on harmonic sums, *Proc. Amer. Math. Soc.* **135** (2007) 1329–1333.

Department of Mathematics, Maritime Faculty, University of Montenegro, Dobrota 36, 85330 Kotor, Montenegro romeo@ac.me